

Scott M. Browning, MD, FACS, FASCRS
Megan M. Cavanaugh, MD, FACS, FASCRS
Jeffrey V. Manchio, MD, FACS, FASCRS
Colon and Rectal Surgeons

OUTPATIENT SURGERY PREP INSTRUCTIONS

LOCATION:	DATE:
	CHECK IN TIME:
SURGERY:	
•	le your procedure, please call us with a minimum of a 2 week notice* ocedure or the above instructions, please contact our office at: 971-254-9884
7 DAYS PRIOR:	<u>_</u>
 coordinate with your prescribing physicia Common medication restrictions: Couma DO NOT TAKE Aspirin, Ibuprofen, Advil, Not any take Tylenol as needed. STOP herbal medicines and supplements 	nning medication you will need to discuss this with our scheduler. We will an on how you should take this medication prior to your surgery. adin (Warfarin), Effient, Eliquis, Plavix (Clopidogrel), Pradaxa, Xarelto. Motrin, Nuprin, Naproxen, Aleve, Naprosyn or NSAIDs. that thin the blood i.e., vitamin E and fish oil. aedications (including blood pressure and heart medications) as usual, th a sip of water. Nothing else.
5 DAYS PRIOR: CONFIRM YOUR RIDE HOME. You may not us	 se a taxi or public transportation unless accompanied by a responsible adult.
SURGERY DAY:	
You may not have anything to eat or drink *This includes: gum, water, candy, coffee, One Enema Prep – Buy this over the co	
rectally, 2 hours before leaving home.	,
☐ Two Enema Prep — Buy this over the corectally, 2 hours before leaving home, s	ounter at your pharmacy and administer per rectum; 2 Fleet enemas spaced 30 minutes apart.
recovery areas. They are welcome to remain	leave all valuables at home. Support persons are welcome in the pre-op and in the waiting room during your procedure. We encourage a support person th you for at least 24 hours after your procedure.
YOUR POST-OP IS ON:	CHECK IN AT:
St. Vincent Office 9155 SW Barnes Rd. Suite 231 Portland, OR 97225	☐ Bridgeport Office 18040 SW Lwr Boones Ferry Rd., Suite 207 Tigard. OR 97224

WHAT TO EXPECT AFTER ANORECTAL SURGERY

- You may pass blood with bowel movements which is normal.
- Call if you pass larger than a "half dollar" sized blood clot or enough blood to fill a coffee cup.
- You may have rectal discharge/drainage/itching or pain for up to 8 weeks after surgery.

DIET AND BOWEL HABITS

- Eat your usual diet with plenty of fruits, vegetables, and bran cereal.
- Avoid spicy food. Drink a minimum of 2 quarts of water daily.
- We strongly recommend a fiber supplement; such as, Citrucel or Metamucil taken once or twice daily.
- Preferably you should have a soft, yet formed bowel movement (BM) daily without straining.
- Your doctor may recommend for you to also take a stool softener; such as, Colace or mineral oil.
- If you have not had a BM within 48 hours, you can take Milk of Magnesia, 1-2 tablespoons every 6-8 hours until you have one.
- If you are having liquid BM's, stop the stool softener.

CARE OF ANUS AND SURROUNDING SKIN

- Following a BM, you may gently wash with a moist Kleenex, squirt bottle (i.e. used shampoo bottle), or a spray bottle with warm water to clean the area.
- Afterward, use a warm bath/Sitz bath for 5-15 minutes to continue cleaning and soothe the area.
- A warm shower will also work, however sometimes tub baths help more. No special additives, to the water, are needed.
- Try to do 3 Sitz baths a day, especially for the first week or two, then less as you feel better.
- Dry thoroughly by patting or using a blow dryer on a warm setting. Feel free to use cotton or gauze between your buttocks to absorb drainage.
- "Donut" cushions are not recommended. Use a soft pad or pillow to sit on instead.

PAIN MEDICATION

- If you need refills that are not included with your original prescription, please call the office during office hours (M-F 9am to 5pm) to request a refill. Refills will not be given on weekends/nights by the on-call physician.
- <u>Allow 3 business days for a refill</u>. Due to Federal law, scheduled pain prescriptions cannot be called in, faxed or electronically prescribed. The majority of post-op pain medications our physicians prescribe fall under this rule.
- Plan on having a friend or family member pick up your prescription from our St. Vincent office.

ACTIVITY

- This is determined by your comfort level.
- Driving should wait until at least 48 hours after surgery. Do not drive within 4 hours of taking pain medications.
- Sexual activity is permitted if it does not cause discomfort. This could be a few days to a few weeks; each patient and each procedure is different.

URINARY ISSUES

- It is very common to experience some difficulty emptying your bladder for the first few days. Warm baths, more activity, and less pain medicine are helpful.
- If you cannot empty your bladder at all and it is getting increasingly painful, go to your nearest ER, because you may need to be catheterized.

FOLLOW-UP APPOINTMENT

- Please plan on attending your scheduled post-op visit.
- If you are experiencing increasing pain, a fever (> 100.6° F), or have questions or concerns, please call the office at 971-254-9884 to be seen sooner.

If you are experiencing a medical emergency, call 911 or go to the emergency room.



Thank you for choosing Providence Milwaukie Hospital to assist you in your care! You are scheduled for a surgery with a doctor from **Colorectal Health NW**. Please read the following instructions to help ease your way.

Before your procedure:

Please register as soon as you know your physician has arranged your procedure. **Call 1-833-386-1128 to register over the phone**. You will be asked to verify your insurance and provide some general information.

Location and Parking:

Providence Milwaukie Hospital is located at 10150 S.E. 32nd Ave Milwaukie, OR 97222. Please take the driveway (off of 32nd Ave) with the Main Entrance/Emergency Entrance sign. Park in the parking lot at the top of the hill on the right side. Go through the Main Entrance sliding glass door. Immediately on your left is the Outpatient Services/Admitting desk, check in there.

Please bring:

- If you wear contact lenses bring your case and solution.
- Your reading glasses, hearing aids, and any mobility devices such as a cane or walker. Wheelchairs are available for use while you are at the hospital.

Please do not bring any valuables or wear any jewelry.

Due to fragrance sensitivity and potential allergic reaction we ask that you and your visitors please do not wear perfume, body spray, cologne or other fragrance while visiting the hospital.

Transportation:

**You MUST have someone drive you home. You CANNOT take a bus or taxi.

Due to the short duration of the procedure, we recommend your driver stay on the hospital premises.

If you have any further questions you may contact your physician's office or contact Pre-Procedural Services at 503-513-8843.

