

INSTRUCTIONS AFTER RUBBER BAND LIGATION

RUBBER BANDS:

There are many methods of treating hemorrhoids. Rubber band ligation is done by application of a small rubber band over the hemorrhoid. The bands, being elastic, tighten and cut off the blood supply to the hemorrhoid until it falls off. Since internal hemorrhoids usually do not have much sensation to pain, this method is easily tolerated. However, external hemorrhoids are very sensitive, and this method cannot be used for them.

AFTER THE PROCEDURE:

It is not uncommon to feel a dull ache for a day or two. This sensation may be noted as soon as the rubber band is applied. Occasionally, it will make you feel as if you want to have a bowel movement. For this ache or discomfort, a great majority of patients either take no medication or they take Tylenol. (NOTE: AVOID THE USE OF ASPIRIN, ADVIL, OR MOTRIN (IBUPROFEN), ALEVE, ARTHRITIS MEDICATIONS, OR BLOOD THINNERS FOR ONE WEEK). Your physician may give you a prescription for a stronger pain medication if needed. Taking warm Sitz baths for 15 minutes will also relieve this discomfort.

DIET:

You may resume your regular diet. It is important to keep your bowel movements reasonably soft. You should take a bulk laxative (bran, Metamucil, Konsyl, Citrucel etc.) daily to avoid hard stools or diarrhea.

ACTIVITY:

You may resume your normal activities such as driving a car, work, and sports.

PRECAUTIONS:

The banded hemorrhoid will drop off in about 3-10 days. Usually you will not notice anything other than some minor bleeding. Should severe bleeding (half a cup of blood at a time or more) occur, you should contact us immediately. If pain persists past 48 hours, or if it is not controlled by pain pills, or if there is urinary difficulty or abdominal pain, please call the office. Although this treatment is safe and effective, rare instances of severe infection or bleeding can occur. The main symptoms of infection are severe pain, chills and fever, and inability to urinate.

FOLLOW-UP TREATMENT:

Generally, it takes between 1-4 treatments at intervals of approximately 2 weeks to eliminate the symptomatic internal hemorrhoid prolapse. Only one treatment is given at a time, in order to minimize pain and complications. Subsequent treatments tend to be less uncomfortable, since they are smaller than the first hemorrhoid treated.

If you have questions, please call the office at 503-216-5380. If you have a medical problem that requires more immediate medical attention, please present to the Emergency Department at St. Vincent Medical Center.